



Periodic Table & Periodicity

Long form of the Periodic Table or Moseley's Periodic Table or Modern Periodic Table :

S.No.	Introduction	DISCRIPTION
1.	Proposed by	Moseley
2.	Contribution	(i) In the long form of periodic table there is contribution of Ramsey, Werner, Bohr and Bury. (ii) This table is also referred to as Bohr's table since it follows Bohr's scheme of the arrangements of elements into four types based on electronic configuration of elements. The modern periodic table consists of horizontal rows (periods) and vertical column (groups).
3.	Based on	Atomic number
4.	Experiment	<p>(i) Moseley did an experiment in which he bombarded high speed electrons on different Metal surfaces and obtained X-rays (electromagnetic rays). He observed regularities in the characteristic X-ray spectra of the elements and found that plot $\sqrt{\nu}$ vs. Z (atomic number) is straight line while $\sqrt{\nu}$ vs. A (atomic weight) is not, and $\sqrt{\nu} = a(Z - b)$, where a and b are constants that are same for all elements and ν is frequency of X-rays. Thus he concluded that atomic number is more fundamental property than atomic weight.</p> <p>(ii) Moseley concluded that the physical and chemical properties of the elements are periodic function of their atomic number.</p>
5.	Modern Periodic Law	The physical and chemical properties of elements are periodic function of their atomic number . So the elements are arranged in order of increasing atomic number, the elements with similar properties comes after regular intervals.
6.	Periodicity	The repetition of the properties of elements after regular intervals when the elements are arranged in the order of increasing atomic number is called periodicity. (a) In a period, the ultimate orbit remain same, but the number of electrons gradually increases. (b) In a group, the number of electrons in the ultimate orbit remains same, but the values of n increases.
7.	Cause of Periodicity	The periodic repetition of the properties of the elements is due to the recurrence of similar valence shell electronic configuration after certain regular intervals. For example, alkali metals have same electronic configuration ns^1 , therefore, have similar properties. In the periodic table, elements with similar properties occur at intervals of 2, 8, 8, 18, 18 and 32. These numbers are called as magic numbers.

The modern periodic table consists of horizontal rows (periods) and vertical column (groups).

Periods: There are seven periods numbered as 1, 2, 3, 4, 5, 6 and 7.

- Each period consists of a series of elements having same valence shell.
- Each period corresponds to a particular principal quantum number of the valence shell present in it.
- Each period starts with an alkali metal having outermost electronic configuration as ns^1 .





Classification of the Elements :

It is based on the type of orbitals which receives the differentiating electron (i.e., last electron).

(a) s-block elements

When shells upto $(n - 1)$ are completely filled and the last electron enters the s-orbital of the outermost (n^{th}) shell, the elements of this class are called s-block elements.

- ◆ Group 1 & 2 elements constitute the s-block.
- ◆ General electronic configuration is [inert gas] ns^{1-2}
- ◆ s-block elements lie on the extreme left of the periodic table.
- ◆ This block includes metals.

(b) p-block elements

When shells upto $(n - 1)$ are completely filled and differentiating electron enters the p-orbital of the n^{th} orbit, elements of this class are called p-block elements.

- ◆ Group 13 to 18 elements constitute the p-block.
- ◆ General electronic configuration is [inert gas] $ns^2 np^{1-6}$
- ◆ p-block elements lie on the extreme right of the periodic table.
- ◆ This block includes some metals, all nonmetals and metalloids.

(c) d-Block elements

When outermost (n^{th}) and penultimate shells $(n - 1)^{\text{th}}$ shells are incompletely filled and differentiating electron enters the $(n - 1)$ d orbitals (i.e., d-orbital of penultimate shell) then elements of this class are called d-block elements.

- ◆ Group 3 to 12 elements constitute the d-block.
- ◆ General electronic configuration is [inert gas] $(n - 1)d^{1-10}ns^{1-2}$ (except, palladium which has valence shell electron configuration $4d^{10}5s^0$).
- ◆ All the transition elements are metals and most of them form coloured complexes or ions.
- ◆ d-block elements are classified into four series as given below.
 - (1) Ist transition series i.e. 3d series contains 10 elements and starts from $_{21}\text{Sc}$ – $_{30}\text{Zn}$. Filling of electrons takes place in 3d sub-shell.
 - (2) IInd transition series i.e. 4d series contains 10 elements and starts from $_{39}\text{Y}$ – $_{48}\text{Cd}$. Filling of electrons takes place in 4d sub-shell.
 - (3) IIIrd transition series i.e. 5d series contains 10 elements and starts from $_{57}\text{La}$, $_{72}\text{Hf}$ – $_{80}\text{Hg}$. Filling of electrons takes place in 5d sub-shell.
 - (4) IVth transition series i.e. 6d series contains 10 elements and starts from $_{89}\text{Ac}$, $_{104}\text{Rf}$ – $_{112}\text{Uub}$. Filling of electrons takes place in 6d sub-shell (incomplete series).
- ◆ Those elements which have partially filled d-orbitals in neutral state or in any stable oxidation state are called transition elements. (Zn, Cd, Hg, Uub are not transition elements)

(d) f-Block elements

When n , $(n - 1)$ and $(n - 2)$ shells are incompletely filled and last electron enters into f-orbital of antepenultimate i.e., $(n - 2)^{\text{th}}$ shell, elements of this class are called f-block elements. General electronic configuration is $(n - 2)f^{1-14}(n - 1)d^{0-1}ns^2$

- ◆ All f-block elements belong to 3rd group.
- ◆ They are metals
- ◆ Within each series, the properties of the elements are quite similar.
- ◆ The elements coming after uranium are called transuranium elements.
- ◆ They are also called as inner-transition elements as they contain three outer most shell incomplete and were also referred to as rare earth elements since their oxides were rare in earlier days.

The elements of f-blocks have been classified into two series.

 1. Ist inner transition or 4 f-series, contains 14 elements $_{58}\text{Ce}$ to $_{71}\text{Lu}$. Filling of electrons takes place in 4f subshell.
 2. IInd inner transition or 5 f-series, contains 14 elements $_{90}\text{Th}$ to $_{103}\text{Lr}$. Filling of electrons takes place in 5f subshell.
- ◆ The actinides and lanthanides have been placed at the bottom of the periodic table to avoid the undue expansion of the periodic table.




Building-Block Elements

Oxygen (O)
Carbon (C)
Hydrogen (H)
Nitrogen (N)

These four elements compose almost 96% of the mass of the human body.

Major Minerals

Potassium (K), sodium (Na), and chlorine (Cl) are present in body fluids.

Magnesium (Mg) and sulfur (S) are found in proteins.


Calcium (Ca) and phosphorus (P) are present in teeth and bones.

Each major mineral is present in 0.1–2% by mass. At least 100 mg of each mineral is needed in the daily diet.

Trace Elements





Arsenic (As)	Iron (Fe)
Boron (B)	Manganese (Mn)
Chromium (Cr)	Molybdenum (Mo)
Cobalt (Co)	Nickel (Ni)
Copper (Cu)	Selenium (Se)
Fluorine (F)	Silicon (Si)
Iodine (I)	Zinc (Zn)

Each trace element is present in less than 0.1% by mass. A small quantity (15 mg or less) of each element is needed in the daily diet.



Solved Examples

Ex-1 Elements A, B, C, D and E have the following electronic configurations :



Which among these will belong to the same group in the periodic table ?

Sol. Out of these, elements A and B will belong to the same group of the periodic table because they have same outer electronic configuration, $ns^2 np^1$.

Prediction of period, group and block:

- Period of an element corresponds to the principal quantum number of the valence shell.
- The block of an element corresponds to the type of subshell which receives the last electron.
- The group is predicted from the number of electrons in the valence shell or/and penultimate shell as follows.
 - (a) For s-block elements, Group number = the number of valence electrons
 - (b) For p-block elements, Group number = 10 + number of valence electrons
 - (c) For d-block elements, Group number = number of electrons in $(n - 1)$ d sub shell + number of electrons in valence shell.
 - (d) For f block element – $\left. \begin{array}{l} (58-71) \quad 4f \\ (92-103) \quad 5f \end{array} \right\}$ - All f block elements are belongs to group number 3.

Two exceptions to this categorization:

Strictly, helium belongs to the s-block but its positioning in the p-block along with other group 18 elements is justified because it has a completely filled valence shell ($1s^2$) and as a result, exhibits properties & characteristics of other noble gases.

Hydrogen has only one s-electron and hence can be placed in group 1 (alkali metals). It can also gain an electron to achieve a noble gas arrangement and hence it can behave similar to a group 17 (halogen family) elements. Because it is a special case, we shall place hydrogen separately at the top of the periodic table.





Isotopes in modern periodic table:

As isotopes of an element have same atomic number and in modern periodic table the elements have been allotted places based on their atomic number. All the isotopes of an element have been assigned the same position in the modern periodic table.

Solved Examples

Ex-2 An element X with $Z = 112$ has been recently discovered. What is the electronic configuration of the element? To which group and period will it belong?

- Sol.**
- The electronic configuration of element X is $[\text{Rn}]^{86} 5f^{14} 6d^{10} 7s^2$
 - It belongs to d-block as last electron enters in d subshell.
 - As number of electrons in $(n - 1)d$ subshell and valence shell is equal to twelve i.e. $10 + 2$. So it belongs to group 12.
 - It belongs to period 7 of the periodic table as principal quantum number of valence shell is 7 (i.e., $7s^2$).

Metals and nonmetals :


- The metals are characterised by their nature of readily giving up the electron(s) and from shining lustre. Metals comprise more than 78% of all known elements and appear on the left hand side of the periodic table. Metals are usually solids at room temperature (except mercury, gallium). They have high melting and boiling points and are good conductors of heat and electricity. Oxides of metals are generally basic in nature (some metals in their higher oxidation state form acid oxides e.g. CrO_3).
- Nonmetals do not lose electrons but take up electrons to form corresponding anions. Nonmetals are located at the top right hand side of the periodic table. Nonmetals are usually solids, liquids or gases at room temperature with low melting and boiling points. They are poor conductors of heat and electricity. Oxides of nonmetals are generally acidic in nature.

Metalloids (Semi metals) :

- It can be understood from the periodic table that nonmetallic character increases as we move from left to right across a row. It has been found that some elements which lie at the border of metallic and nonmetallic behavior, possess the properties that are characteristic of both metals and nonmetals. These elements are called semi metals or metalloids.
- The metalloids comprise of the elements Si, Ge, As, Sb, Se and Te.
- Oxides of metalloids are generally amphoteric in nature.

Diagonal relationship :

Some elements of certain groups of 2nd period resemble much in properties with the elements of third period of next group i.e. elements of second and third period are diagonally related in properties. This phenomenon is known as diagonal relationship. For example, the similarity between lithium (the first member of group 1) and magnesium (the second element in group 2) is called a diagonal relationship. Diagonal relationship also exist between other pairs of elements Be and Al, B and Si as shown in figure;



Diagonal relationship arises because of ;

- on descending a group, the atoms and ions increase in size. On moving from left to right in the periodic table, the size decreases. Thus on moving diagonally, the size remains nearly the same. ($\text{Li} = 1.23 \text{ \AA}$ & $\text{Mg} = 1.36 \text{ \AA}$; $\text{Li}^+ = 0.76 \text{ \AA}$ & $\text{Mg}^{2+} = 0.72 \text{ \AA}$)
- it is sometimes suggested that the diagonal relationship arises because of diagonal similarity in electronegativity values. ($\text{Li} = 1.0$ & $\text{Mg} = 1.2$; $\text{Be} = 1.5$ & $\text{Al} = 1.5$; $\text{B} = 2.0$ & $\text{Si} = 1.8$)
- Be and Al also show a diagonal relationship. In this case sizes are not so close ($\text{Be}^{2+} = 0.45 \text{ \AA}$ and $\text{Al}^{3+} = 0.535 \text{ \AA}$) but the charge / ionic radius ratio is nearly similar because the charges are 2+ and 3+ respectively.

$$\text{Charge / Ionic radius ratio} = \frac{(\text{Ionic charge})}{(\text{Ionic radius})}$$



Shielding Effect and Z_{eff}

The periodicity of atomic properties :

(1) Effective nuclear charge (Z_{eff}):

The outer electron experiences two opposing force :-

- (i) Force of attraction from nucleus.
- (ii) Force of repulsion from inner electrons.

Suppose inner electrons are not present there then force of attraction experience by test electron must be greater than it actually experienced because electronic repulsion weakens the force of attraction.

This reduced nuclear charge or nuclear actually experienced by an electron is termed as effective nuclear charge.

$$Z_{\text{eff}} = Z - \sigma$$

Z_{eff} = Effective nuclear charge

Z = Atomic number

σ = Screening constant or shielding constant or Slater constant

Effective nuclear charge is not same for all the electrons present in an atom and it varies with distance between electron and nucleus.



This effect in which inner electrons shield valence electron from the attraction due to nucleus is called shielding effect.

It is also known as screening effect as inner electron create a protective screen between nucleus and valence electrons.

Due to shielding effect valence electron experiences less attraction from the side of nucleus as if inner electrons are absent.


s-orbitals have the largest screening effect for a given n value since s electrons are closer to the nucleus.

Screening effect decreases as s-orbital > p-orbital > d-orbital > f-orbital.




Section (A) : Atomic and Ionic radius
(1) Atomic radius :

It is distance between outermost electron and nucleus.



X-ray diffraction, electron diffraction method and nuclear magnetic resonance (NMR) spectrum methods are used to determine internuclear distance or bond length.

Atomic radius depends on the type of chemical bond between atoms in a molecule. These are :

1. Covalent radius
2. Ionic radius
3. Metallic radius
4. Vander waal's radius

S.No.	Type of radius	DISCRIPTION				
1.	Covalent radius SBCR(Single Bonded Covalent Radius)	<p>Covalent radius is calculated in covalent compounds. Covalent compound's form covalent bonds. Covalent bonds are formed by overlapping of atomic orbitals. Type of Covalent radius.</p>				
	Halogen	Molecule	Structure	Model	$d(X-X)$ / pm (gas phase)	$d(X-X)$ / pm (solid phase)
	Fluorine	F ₂	$\begin{array}{c} \text{F} - \text{F} \\ \longleftrightarrow \\ 143\text{pm} \end{array}$		143	149
	Chlorine	Cl ₂	$\begin{array}{c} \text{Cl} - \text{Cl} \\ \longleftrightarrow \\ 199\text{pm} \end{array}$		199	198
	Bromine	Br ₂	$\begin{array}{c} \text{Br} - \text{Br} \\ \longleftrightarrow \\ 228\text{pm} \end{array}$		228	227
	Iodine	I ₂	$\begin{array}{c} \text{I} - \text{I} \\ \longleftrightarrow \\ 266\text{pm} \end{array}$		266	272
	(i) In Homo	Covalent radius is the one-half of the internuclear distance between two singly				



	atomic Molecules	bonded two nuclei (of like atoms). Internuclear distance of A–A(A ₂) molecule is (d _{A–A}) and covalent radius is r _A then $d_{A-A} = r_A + r_A \Rightarrow d_{A-A} = 2r_A$ $r_A = \frac{d_{A-A}}{2}$												
		e.g. – In Cl ₂ molecule, internuclear distance is 1.98 Å so $r_{Cl} = \frac{1.98}{2} = 0.99 \text{ Å}$												
		SBCR of O, N and C etc. elements can be determined by taking H ₂ O ₂ , N ₂ H ₄ , C ₂ H ₆ respectively.												
	(ii) Hetero atomic molecule	(a) For Hetero atomic molecule with not E.N. difference. $d_{A-B} = r_A + r_B$												
		Example : For A–B molecule Electronegativities of A and B are approximately equal e.g. C–I E.N. of C and I are approx equal (2.5) internuclear distance of C–I is 2.13Å and r _C is 0.77Å. Solution : d _{C–I} = r _C + r _I ∴ r _I = 2.13 – 0.77 = 1.36Å												
		(b) Heteroatomic molecule with Δ E.N. difference more If in a diatomic molecule electronegativities of A–B have more difference. Then actual bond length will be reduced.												
2.	Ionic Radius													
	(i) Cationic radius	When an neutral gaseous atom loses electron it converts into cation. Radius = Atomic radius > Cationic radius Reason: After loosing electron number of electron reduces, but number of protons remains same, due to this Z _{eff} increases, hence electrons pulls towards nucleus and atomic radius decreases, moreover after losing all the electrons from the outer most shell, penultimate shell becomes ultimate shell which is nearer to nucleus so size decreases.												
		$\text{Size of cation} \propto \frac{1}{\text{Magnitude of +ve charge or } Z_{\text{eff}}}$												
		Example $\text{Size of cation} = \text{Fe} > \text{Fe}^{+2} > \text{Fe}^{+3}$ $\text{Size of cation} = \text{Pb}^{+2} > \text{Pb}^{+4}$ $\text{Size of cation} = \text{Mn} > \text{Mn}^{+2} > \text{Mn}^{+3} > \text{Mn}^{+4} > \text{Mn}^{+5} > \text{Mn}^{+6} > \text{Mn}^{+7}$												
	(ii) Anionic radius	When neutral gaseous atom gains electron it converts into anion. Radius = Anionic radius > Atomic radius Reason: In an anion electron are more than protons so effective nuclear charge reduces, and inter electronic repulsion increases, which also increases screening effect. So distance between electron and nucleus increases and size of anion also increases.												
		$\text{Size of anion} \propto \text{Magnitude of -ve charge}$												
	radius	Example : Size of F ⁻ > F why ? Sol. <table border="1" style="margin-left: auto; margin-right: auto;"> <tbody> <tr> <td></td> <td>F</td> <td>F⁻</td> </tr> <tr> <td>Proton</td> <td>9</td> <td>9</td> </tr> <tr> <td>electron</td> <td>9</td> <td>10</td> </tr> <tr> <td>$\frac{Z}{e^{\ominus}}$</td> <td>$\frac{9}{9} = 1$</td> <td>$\frac{9}{10} = 0.9$</td> </tr> </tbody> </table> As Z _{eff} of F ⁻ is less than F so size of F ⁻ > F		F	F ⁻	Proton	9	9	electron	9	10	$\frac{Z}{e^{\ominus}}$	$\frac{9}{9} = 1$	$\frac{9}{10} = 0.9$
	F	F ⁻												
Proton	9	9												
electron	9	10												
$\frac{Z}{e^{\ominus}}$	$\frac{9}{9} = 1$	$\frac{9}{10} = 0.9$												





Sizes of atoms and their ions in pm

Group 1		Group 2		Group 13		Group 16		Group 17	
Li ⁺ 90	Li 134	Be ²⁺ 59	Be 90	B ³⁺ 41	B 82	O 73	O ²⁻ 126	F 71	F ⁻ 119
Na ⁺ 116	Na 154	Mg ²⁺ 86	Mg 130	Al ³⁺ 68	Al 118	S 102	S ²⁻ 170	Cl 99	Cl ⁻ 167
K ⁺ 152	K 196	Ca ²⁺ 114	Ca 174	Ga ³⁺ 76	Ga 126	Se 116	Se ²⁻ 184	Br 114	Br ⁻ 182
Rb ⁺ 166	Rb 211	Sr ²⁺ 132	Sr 192	In ³⁺ 94	In 144	Te 135	Te ²⁻ 207	I 133	I ⁻ 206

(iii) Size of iso electronic species

					H ⁻	He
Li ⁺	Be ²⁺		N ³⁻	O ²⁻	F ⁻	Ne
Na ⁺	Mg ²⁺	Al ³⁺		S ²⁻	Cl ⁻	Ar
K ⁺	Ca ²⁺	Sc ³⁺		Se ²⁻	Br ⁻	Kr
Rb ⁺	Sr ²⁺	Y ³⁺		Te ²⁻	I ⁻	Xe
Cs ⁺	Ba ²⁺	La ³⁺				

Those species having same number of electron but different nuclear charge forms isoelectronic series.
For isoelectronic species the atomic radius increases with decrease in nuclear charge.

Ions	K ⁺	Ca ²⁺	S ²⁻	Cl ⁻
Atomic number	19	20	16	17
electron	18	18	18	18

Example


Order of radius = S²⁻ > Cl⁻ > K⁺ > Ca²⁺

Order of radius = P³⁻ > S²⁻ > Cl⁻

Order of radius = N³⁻ > O²⁻ > F⁻ > Na⁺ > Mg²⁺ > Al³⁺

Order of radius = Au⁺ > Hg²⁺ > Tl³⁺ > Pb⁴⁺

3. **Metallic radius**



It is one-half of the nuclear distance between two adjacent metallic atoms in crystalline lattice structure.
so metallic radius > Covalent radius.

$$\text{Metallic radius} \propto \frac{1}{\text{Metallic bond strength}}$$

More metallic radius → loose crystal packing → less bond strength (body centered packing)

Less metallic radius → tight crystal packing FCC → High bond strength (Hexagonal close packing)



4.	Van der Waal's radius	Those atoms (like noble gases) which are not bonded with each other, experiences a weak attractive force to come nearer.	
		The half of the distance between the nuclei of adjacently placed atoms in solid state of a noble gas is Vander Waal's radius.	
		Vander Waal radius > Covalent radius.	
		Inert gas have only Vander Waal radius	
		In molecules of nonmetals both covalent and Vander Waal radius exists.	
Radius = Vander Waal radius > Metallic Radius > Covalent Radius			
Van der waal force of attraction \propto Molecular weight OR atomic weight (in inert gases)			
In a period from left to right Vander Waal radius decreases. In a group from top to bottom its values increases.			
5.	Factors affecting atomic size		
	(i) Effective nuclear charge	Atomic radius $\propto \frac{1}{\text{Effective nuclear charge}}$	Example Li > Be > B > C > N > O > F
	(ii) Number of shells	Atomic radius \propto No. of shells	Example Li < Na < K < Rb < Cs
	(iii) screening effect	Atomic radius \propto Screening effect	
	(iv) Magnitude of +ve charge	Atomic radius $\propto \frac{1}{\text{Magnitude of +ve charge}}$	Example Mn > Mn ⁺² > Mn ⁺³ > Mn ⁺⁴
	(v) Magnitude of -ve charge	Atomic radius \propto Magnitude of -ve charge	Example O < O ⁻ < O ⁻²
	(vi) Bond order	Atomic radius $\propto \frac{1}{\text{Bond order}}$	Example N-N > N=N > N≡N




Table-3

Variation in a Period (left to right) :	Variation in a Group (top to bottom) :
Nuclear charge (Z) increases by one unit	Nuclear charge (Z) increases by more than one unit
Effective nuclear charge (Z_{eff}) also increases	Effective nuclear charge (Z_{eff}) almost remains constant because of increased screening effect of inner shells electrons.
But number of orbitals (n) remains constant	But number of orbitals (n) increases.
As a result, the electrons are pulled closer to the nucleus by the increased Z_{eff} . $r_n \propto \frac{1}{Z^*}$ Hence atomic radii decrease with increase in atomic number in a period from left to right.	The effect of increased number of atomic shells overweighs the effect of increased nuclear charge. As a result of this the size of atom increases from top to bottom in a given group.

Irregularities

- In the transition series (e.g. in first transition series), the covalent radii of the elements decrease from left to right across a row until near the end when the size increases slightly. On passing from left to right, extra protons are placed in the nucleus and the extra electrons are added. The orbital electrons shield the nuclear charge incompletely. Thus the nuclear charge attracts all the electrons more strongly, hence a contraction in size occurs. The radii of the elements from Cr to Cu, are very close to one another because the successive addition of d-electrons screen the outer electrons (4s) from the inward pull of the nucleus. As a result of this, the size of the atom does not change much in moving from Cr to Cu.

Table-4

Element	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
Atomic radius (pm)	144	132	122	117	117	117	116	115	117	125

- The elements of group 3 belonging to d-block show the expected increase in size as that found in case of s-block and p-block elements. However, in the subsequent groups that is 4th onwards (upto 12th group), there is increase in size between the first and second member, but hardly any increase between the second and third elements of the given group. There are 14 lanthanide elements between lanthanum and hafnium, in which the antepenultimate 4f shell of electrons (exert very poor shielding effect) is filled. There is a gradual decrease in size of the 14 lanthanide elements from cerium to lutetium. This is called lanthanide contraction. This lanthanide contraction cancels out the normal size increase on descending a group in case of transition elements.

Solved Examples

Ex-3 Atomic radius of Li is 1.23 Å and ionic radius of Li^+ is 0.76 Å. Calculate the percentage of volume occupied by single valence electron in Li.

Sol. Volume of Li = $\frac{4}{3} \times 3.14 \times (1.23)^3 = 7.79 \text{ Å}^3$ ($-\text{Li} = 1s^2 2s^1$)

Volume of Li^+ = $\frac{4}{3} \times 3.14 \times (0.76)^3 = 1.84 \text{ Å}^3$ ($-\text{Li}^+ = 1s^2$)

∴ Volume occupied by 2s subshell = $7.79 - 1.84 = 5.95 \text{ Å}^3$.

∴ % Volume occupied by single valence electron i.e., 2s electron = $\frac{5.95}{7.79} \times 100 = 76.4\%$

Ex-4 Select from each group the species which has the smallest radius stating appropriate reason.

(a) O, O^- , O^{2-} (b) P^{3+} , P^{4+} , P^{5+}

Sol. (a) O is having smallest radius. Anion is larger than its parent atom. Also the anion of the same atom with higher negative charge is bigger in size as compared to anion with smaller negative charge as proton to electron ratio decreases thus attraction between valence shell electrons and nucleus decreases. Hence electron cloud expands.





(b) The ionic radius decreases as more electrons are ionized off that is as the valency increases. So the correct order is $P^{5+} < P^{4+} < P^{3+}$.


Ex-5 Mg^{2+} is smaller than O^{2-} in size, though both have same electronic configuration. Explain ?

Sol. Mg^{2+} and O^{2-} both are isoelectronic i.e., have same number of electrons. But Mg^{2+} having 12 protons in its nucleus exerts higher effective nuclear charge than O^{2-} having 8 protons and thus valence shell as well as inner shells electrons are more strongly attracted by the nucleus in Mg^{2+} resulting smaller size than O^{2-} .

Section (B) : Ionisation energy

(2) Ionisation Energy :

Ionisation energy (IE) is defined as the amount of energy required to remove the most loosely bound electron from an isolated gaseous atom to form a cation.



IE_1 , IE_2 & IE_3 are the 1st, 2nd & 3rd ionization energies to remove electron from a neutral atom, monovalent and divalent cations respectively.

In general, $(IE)_1 < (IE)_2 < (IE)_3 < \dots$ because, as the number of electrons decreases, the attraction between the nucleus and the remaining electrons increases considerably and hence subsequent ionization energies increase.

♦ Ionization energies are determined from spectra and are measured in kJ mol^{-1} , k Cal mol^{-1} , eV (electron volt).

♦ **Factors Influencing Ionisation energy**

Variation in ionization energies in a period and group may or not be regular and can be influenced by the following factors.

(A) Size of the Atom : Ionisation energy decreases with increase in atomic size.

As the distance between the outer most electrons and the nucleus increases, the force of attraction between the valence shell electrons and the nucleus decreases. As a result, outer most electrons are held less firmly and lesser amount of energy is required to knock them out.

For example, ionisation energy decreases continuously in a group from top to bottom with increase in atomic size. But in 13th & 14th group the ionisation energy does not decrease continuously and order is as follows:

For 13th group: $B > Tl > Ga > Al > In$


For 14th group: $C > Si > Ge > Pb > Sn$

(B) Nuclear Charge : The ionisation energy increases with increase in the nuclear charge.

This is due to the fact that with increase in the nuclear charge, the electrons of the outer most shell are more firmly held by the nucleus and thus greater amount of energy is required to pull out an electron from the atom.

For example, ionisation energy increases as we move from left to right along a period due to increase in nuclear charge.

(C) Shielding or screening effect : The electrons in the inner shells act as a screen or shield between the nucleus and the electrons in the outer most shell. This is called shielding effect. The larger the number of electrons in the inner shells, greater is the screening effect and smaller the force of attraction and thus ionization energy (IE) decreases.



(D) Penetration effect of the electron : The ionization energy also depends on the type of electron which is removed. s, p, d and f electrons have orbitals with different shapes. An s electron penetrates closer to the nucleus, and is therefore more tightly held than a p electron. Similarly p-orbital electron is more tightly held than a d-orbital electron and a d-orbital electron is more tightly held than an f-orbital electron. If other factors being equal, ionisation energies are in the order $s > p > d > f$.


For example, ionisation energy of aluminium is comparatively less than magnesium because outer most electron is to be removed from 3p-orbital (having lesser penetration effect) in aluminium where as in magnesium it will be removed from 3s-orbital (having larger penetration effect) of same energy level.




(E) Electronic Configuration:

If an atom has exactly half-filled or completely filled orbitals, then such an arrangement has extra stability.

The removal of an electron from such an atom requires more energy than expected. For example, first ionisation energy of beryllium is greater than boron because beryllium has extra stable outer most completely filled outer most 2s orbital while boron has partially filled less stable outer most 2p-orbital.



Similarly noble gases have completely filled electronic configurations and hence they have highest ionisation energies in their respective periods.

- Metallic or electropositive character of elements increases as the value of ionisation energy decreases.
- The relative reactivity of the metals in gaseous phase increases with the decrease in ionisation energy.
- The reducing power of elements in gaseous phase increases as the value of ionisation energy decreases. Amongst alkali metals, the lithium is strongest reducing agent in aqueous solution.

Solved Examples


Ex-6 From each set, choose the atom which has the largest ionization enthalpy and explain your answer with suitable reasons. (a) F, O, N (b) Mg, P, Ar

Sol. (a) Fluorine (F) has the largest ionization enthalpy because in moving from left to right in a period, atomic size decreases and electrons are held more tightly. Since F has the smallest size and maximum nuclear charge. It has the largest ionization enthalpy among these elements.
(b) Argon (Ar) has the largest ionization enthalpy as argon has extra stable fully filled configuration.

Ex-7 First and second ionisation energies of magnesium are 7.646 eV and 15.035 eV respectively. The amount of energy in kJ needed to convert all the atoms of magnesium into Mg^{2+} ions present in 12 mg of magnesium vapour will be ? [Given 1 eV = 96.5 kJ mol⁻¹].

Sol. Total energy needed to convert one Mg atom into Mg^{2+} gas ion,
 $= \text{IE}_I + \text{IE}_{II} = 22.681 \text{ eV} = 2188.6 \text{ kJ mol}^{-1}$.
 \Rightarrow 12 mg of Mg = 0.5×10^{-3} mole.
 \therefore Total energy = $0.5 \times 10^{-3} \times 2188.6 = 1.0943 \text{ kJ}$ **Ans.**

Ex-8 Following graph shows variation of ionization energies with atomic number in second period (Li–Ne). Value of ionization energies of Na(11) will be :



- (A) above Ne (B) below Ne but above O.
 (C) below Li (D) between N and O.

Sol. Na is 3rd period element and is bigger than Li. The distance between the nucleus and outer most electron is more as compared to Li. Thus the outer most electron is loosely bound with nucleus and removal of electron is easier. So option (C) is correct.

Ex-9 $\text{M(g)} \longrightarrow \text{M}^+(\text{g}) + \text{e}^-$; $\Delta H = 100 \text{ eV}$.
 $\text{M(g)} \longrightarrow \text{M}^{2+}(\text{g}) + 2\text{e}^-$; $\Delta H = 250 \text{ eV}$.

Which is/are correct statement(s) ?

- (A) IE_1 of M(g) is 100 eV (B) IE_1 of $\text{M}^+(\text{g})$ is 150 eV.
 (C) IE_2 of M(g) is 250 eV. (D) IE_2 of M(g) is 150 eV.

Sol. $\text{M(g)} \longrightarrow \text{M}^+(\text{g}) + \text{e}^-$ IE_1 of M
 $\text{M}^+ \longrightarrow \text{M}^{2+} + \text{e}^-$ IE_2 of M but IE_1 of M^+
 $\text{M} \longrightarrow \text{M}^{2+} + 2\text{e}^-$ ($\text{IE}_1 + \text{IE}_2$). **Ans. (A,B,D)**







Section (C) : Electron gain enthalpy

(3) Electron Affinity:

Electron affinity is conventionally defined as the energy released when an electron is added to the valence shell of an isolated gaseous atom.




A positive electron affinity indicates that the ion X^{-} has a lower, more negative energy than the neutral atom X . The second electron gain enthalpy, the enthalpy change for the addition of a second electron to an initially neutral atom, invariably positive because the electron repulsion outweighs the nuclear attraction.



An element has a high electron affinity if the additional electron can enter a shell where it experiences a strong effective nuclear charge.

The electron gain enthalpy $\Delta_{eg}H^{\ominus}$, is the change in standard molar enthalpy when a neutral gaseous atom gains an electron to form an anion.



Now a days both electron affinity and electron gain enthalpy terms are used.

Both are same in magnitude but opposite in sign.

$$E.A. = +ve \text{ (Exothermic)}: -\Delta_{eg}H^{\ominus} < 0.$$

$$E.A. = -ve \text{ (Endothermic)}: -\Delta_{eg}H^{\ominus} > 0.$$

The units of EA and Electron gain enthalpy are eV/atom or Kcal / mole or KJ / Mole.

Electron gain enthalpy provides a measure of the ease with which an atom adds an electron to form an anion. Electron gain may be either exothermic or endothermic depending on the elements.

When an electron is added to the atom and the energy is released, the electron gain enthalpy is negative and when energy is needed to add an electron to the atom, the electron gain enthalpy is positive.

Although the electron gain enthalpy is the thermodynamically appropriate term, much of inorganic chemistry is discussed in terms of a closely related property, the electron affinity, E_a of an element which is the difference in energy between the gaseous atoms and the gaseous ions.

$$E_a = E(X, g) - E(X^{-}, g)$$

- Group 17 elements (halogens) have very high negative electron gain enthalpies (i.e. high electron affinity) because they can attain stable noble gas electronic configuration by picking up an electron.
- Across a period, with increase in atomic number, electron gain enthalpy becomes more negative because left to right across a period effective nuclear charge increases and consequently it will be easier to add an electron to a small atom.
- As we move in a group from top to bottom, electron gain enthalpy becomes less negative because the size of the atom increases and the added electron would be at larger distance from the nucleus.

In general the electron gain enthalpies for some third period element (e.g. P, S, Cl) are more negative than the corresponding second period members (e.g. N, O, F). This is due to the smaller size of the atoms of the second period elements which would produce larger electron-electron repulsions for the additional electron.

Group 1	$\Delta_{e.g.} H$	Group 16	$\Delta_{e.g.} H$	Group 17	$\Delta_{e.g.} H$	Group 0	$\Delta_{e.g.} H$
H	-73					He	+48
Li	-60	O	-141	F	-328	Ne	+116
Na	-53	S	-200	Cl	-349	Ar	+96
K	-48	Se	-195	Br	-325	Kr	+96
Rb	-47	Te	-190	I	-295	Xe	+77
Cs	-46	Po	-174	At	-270	Rn	+68

- Noble gases have large positive electron gain enthalpies because the electron has to enter the next higher energy level leading to a very unstable electronic configuration.



- Negative electron gain enthalpy of O or F is less than S or Cl. This is due to the fact that when an electron is added to O or F, the added electron goes to the smaller $n = 2$ energy level and experiences significant repulsion from the other electrons present in this level. In S or Cl, the electron goes to the larger $n = 3$ energy level and consequently occupies a larger region of space leading to much less electron-electron repulsion.
- Electron gain enthalpies of alkaline earth metals are very less or positive because the extra electron is to be added to completely filled s-orbitals in their valence shells.
- Nitrogen has very low electron affinity because there is high electron repulsion when the incoming electron enters an orbital that is already half filled.
- (i) Electron affinity $\propto \frac{1}{\text{Atomic size}}$
- (ii) Electron affinity \propto Effective nuclear charge (Z_{eff})
- (iii) Electron affinity $\propto \frac{1}{\text{Screening effect}}$
- (iv) Stability of half filled and completely filled orbitals of a subshell is comparatively more and the addition of an extra electron to such an system is difficult and hence the electron affinity value decreases.

Solved Examples

Ex-10 Consider the elements N, P, O and S and arrange them in order of increasing negative electron gain enthalpy.

Sol. Order of increasing negative electron gain enthalpy is $N < P < O < S$. For detail refer text.

Ex-11 Why do halogens have high electron gain enthalpies (i.e. $-\Delta_{\text{eg}}H^{\ominus}$) ?

Sol. The valence shell electronic configuration of halogens is ns^2np^5 and thus they require one electron to acquire the stable noble gas configuration ns^2np^6 . Because of this they have strong tendency to accept an additional electron and hence have high electron gain enthalpies.

Ex-12 Which will have the maximum value of electron affinity O^x, O^y, O^z [x, y and z respectively are 0, -1 and -2] ?

(A) O^x (B) O^y (C) O^z (D) All have equal.

Sol. Being neutral atom oxygen will have higher electron affinity as there is electrostatic repulsion between additional electron and negative ion in case of O^- and O^{2-} . So option (A) is correct.

Ex-13 The amount of energy when million atoms of iodine are completely converted into I^- ions in the vapour state according to the equation, $I(g) + e^-(g) \rightarrow I^-(g)$ is 5.0×10^{-13} J.

Calculate the electron gain enthalpy of iodine in terms of kJ mol^{-1} and eV per atom.

Sol. The electron gain enthalpy of iodine is equal to the energy released when 1 mole of iodine atoms in vapour state are converted into I^- ions.

$$= -\frac{5.0 \times 10^{-13} \times 6.023 \times 10^{23}}{10^6} = -30.1 \times 10^4 \text{ J} = -301 \text{ kJ.}$$

$$\text{Electron gain enthalpy of iodine in eV per atom} = \frac{-301}{96.5} = -3.12.$$

Ex-14 Account for the large decrease in electron affinity between Li and Be despite the increase in nuclear charge.

Sol. The electron configurations of Li and Be are $[\text{He}]2s^1$ and $[\text{He}]2s^2$, respectively. The additional electron enters the 2s orbital of Li but the 2p orbital of Be and hence is much less tightly bound. In fact, the nuclear charge is so well shielded in Be that electron gain is endothermic.




Section (D) : Electronegativity
(4) Electronegativity:

Electronegativity is a measure of the tendency of an element to attract shared electrons towards itself in a covalently bonded molecules.

The magnitude of electronegativity of an element depends upon its ionisation potential & electron affinity. Higher ionisation potential & electron affinity values indicate higher electronegativity value.

- With increase in atomic size the distance between nucleus and valence shell electrons increases, therefore, the force of attraction between the nucleus and the valence shell electrons decreases and hence the electronegativity values also decrease.
- With increase in nuclear charge force of attraction between nucleus and the valence shell electrons increases and, therefore, electronegativity value increases
- In higher oxidation state, the element has higher magnitude of positive charge. Thus, due to more positive charge on element, it has higher polarising power. Hence with increase in the oxidation state of element, its electronegativity also increases.
- Charge on cation \propto electronegativity of the atom.
- The electronegativity also increases as the s-character in the hybrid orbitals increases.

Hybrid orbital	sp ³	sp ²	sp
s-character	25%	33%	50%

Electronegativity increases \rightarrow

Electronegativity of some elements according to Pauling scale.

Elements	H	Li	Be	B	C	N	O	F	Ne	P	S	Cl	Br	I
Electronegativity	2.1	1.0	1.5	2.0	2.5	3.0	3.5	4.0	0.9	2.1	2.5	3.0	2.8	2.5

Table-5

Variation of electronegativity in a group	Variation of electronegativity in a period
On moving down the groups, Z increases but Z _{eff} almost remains constant, number of shells (n) increases, r _n (atomic radius) increases. Therefore, electronegativity decreases moving down the groups.	While moving across a period left to right, Z, Z _{eff} increases & r _n decreases. Therefore, electronegativity increases along a period.

There is no direct method to measure the value of electronegativity, however, there are some scales to measure its value.

- (a) Pauling's scale :** Linus Pauling developed a method for calculating relative electronegativities of most elements. According to Pauling

$$\Delta = X_A - X_B = 0.208 \sqrt{E_{A-B} - \sqrt{E_{A-A} \times E_{B-B}}}^*$$

E_{A-B} = Bond enthalpy/ Bond energy of A – B bond.

E_{A-A} = Bond energy of A – A bond

E_{B-B} = Bond energy of B – B bond

(All bond energies are in kcal / mol)

$$\Delta = X_A - X_B = 0.1017 \sqrt{E_{A-B} - \sqrt{E_{A-A} \times E_{B-B}}}^* \quad \text{All bond energies are in kJ / mol.}$$

- (b) Mulliken's scale :** Electronegativity χ (chi) can be regarded as the average of the ionisation energy (IE) and the electron affinity (EA) of an atom (both expressed in electron volts).

$$\chi_M = \frac{IE + EA}{2}$$

[Pauling's electronegativity χ_P is related to Mulliken's electronegativity χ_M as given below.

$$\chi_P = 1.35 (\chi_M)^{1/2} - 1.37$$

Mulliken's values were about 2.8 times larger than the Pauling's values.]*

* Only for reference, student need not memorize it.





APPLICATION OF ELECTRONEGATIVITY :

(a) **Nomenclature :**

Compounds formed from two nonmetals are called binary compounds. Name of more electronegative element is written at the end and 'ide' is suffixed to it. The name of less electronegative element is written before the name of more electronegative element of the formula. For example -

Correct formula	Name
(a) I ⁺ Cl ⁻	Iodine chloride
(b) Cl ⁺ F ⁻	Chlorine fluoride
(c) Br ⁺ Cl ⁻	Bromine chloride
(d) IBr	Iodine bromide
(e) OF ₂	Oxygen difluoride
(f) Cl ₂ O	Dichlorine oxide

(b) **Effect on bond length:** When the Δ EN increases between two atoms in a molecule then bond length between that atoms decreases.

Order of Δ EN: - HF > HCl > HBr > HI

Order of bond length: - HF < HCl < HBr < HI

